

The one-dimensional XXZ model with long-range interactions

L. L. Gonçalves¹, A. P. Vieira² and L. P. S. Coutinho¹

February 1, 2008

Abstract

The one-dimensional XXZ model ($s = 1/2$, N sites) with uniform long-range interactions among the transverse components of the spins is considered. The Hamiltonian of the model is explicitly given by $H = J \sum_{j=1}^N (s_j^x s_{j+1}^x + s_j^y s_{j+1}^y) - (I/N) \sum_{j,k=1}^N s_j^z s_k^z - h \sum_{j=1}^N s_j^z$, where the $s^{x,y,z}$ are half the Pauli spin matrices. The model is exactly solved by applying the Jordan-Wigner fermionization, followed by a Gaussian transformation. In the absence of the long-range interactions ($I = 0$), the model, which reduces to the isotropic XY model, is known to exhibit a second-order quantum phase transition driven by the field at zero temperature. It is shown that in the presence of the long-range interactions ($I \neq 0$) the nature of the transition is strongly affected. For $I > 0$, which favours the ordering of the transverse components of the spins, the transition is changed from second- to first-order, due to the competition between transverse and xy couplings. On the other hand, for $I < 0$, which induces complete frustration of the spins, a second-order transition is still present, although the system is driven out of its usual universality class, and its critical exponents assume typical mean-field values.

¹Departamento de Física da UFC, Campus do Pici, Cx. Postal 6030, 60451-970 Fortaleza (CE), Brazil

²Instituto de Física, Universidade de São Paulo, Cx. Postal 66318, 05315-970 São Paulo (SP), Brazil

The observed critical behaviour of magnetic materials, in the very low temperature limit, has renewed the interest in the study of magnetic quantum transitions (S.L.Sondhi,S.M.Girvin,J.P.Carini and D.Shahar, Rev. Mod. Phys. **69**, 315(1997)) . Since these transitions, which are governed by quantum fluctuations, occur at $T=0$, the one-dimensional models play a important role in their study . Therefore, we will consider the exactly soluble one-dimensional XXZ model ($s = 1/2$), with a uniform long-range interaction among the spins along the z direction. Due to the long-range interaction, the model also presents classical critical behaviour, with transitions of first and second order, and it has already been considered by Suzuki (J. Phys. Soc. Jpn. **21**, 2140 (1966)). Since his study was restricted to the analysis of the classical second order transition of the model, and we are interested in its quantum transitions, the model will be considered again. In particular, we will be interested in the effect of the long-range interaction on its quantum critical behaviour.

The Hamiltonian of the model is given by

$$H = J \sum_{j=1}^N (s_j^x s_{j+1}^x + s_j^y s_{j+1}^y) - \frac{I}{N} \sum_{j,k=1}^N s_j^z s_k^z - h \sum_{j=1}^N s_j^z, \quad (1)$$

where $J > 0$, N is the number of sites of the lattice and we assume periodic boundary conditions. By applying the Jordan-Wigner fermionization[1, 2], followed by a Gaussian transformation, we can write the partition function of the model as

$$Z_N = \text{Tr} e^{-\beta H} = C(\beta) \int e^{-\frac{N}{2} z^2} \text{Tr} e^{-\tilde{H}(z)} dz, \quad (2)$$

with

$$\tilde{H}(z) = \frac{\beta J}{2} \sum_{j=1}^N (c_j^\dagger c_{j+1} + c_{j+1}^\dagger c_j) - \tilde{h}(z) \sum_{j=1}^N c_j^\dagger c_j, \quad (3)$$

where $\tilde{h}(z) = \beta(h - I) + \sqrt{2\beta I}z$, $C(\beta)$ depends only on the temperature, a boundary term has been neglected in $\tilde{H}(z)$, and the c_j are fermion operators.

Introducing the Fourier transforms

$$c_j = \frac{1}{\sqrt{N}} \sum e^{-ikj} \hat{c}_k, \quad (4)$$

we can rewrite $\tilde{H}(z)$ in the diagonal form

$$\tilde{H}(z) = \sum_k \tilde{\varepsilon}_k(z) \hat{c}_k^\dagger \hat{c}_k, \quad (5)$$

where $\tilde{\varepsilon}_k(z) = \beta J \cos k - \tilde{h}(z)$ and, due to the periodic boundary conditions, $k = 2\pi n/N$ ($n = 1, \dots, N$). The partition function is then given by

$$Z_N = C(\beta) \int e^{-\frac{N}{2}z^2} \prod_k [1 + e^{-\tilde{\varepsilon}_k(z)}] dz, \quad (6)$$

which, in the thermodynamic limit ($N \rightarrow \infty$), can be evaluated by the saddle-point method. By explicit calculation, we conclude that

$$m \equiv \left\langle \frac{1}{N} \sum_{j=1}^N s_j^z \right\rangle = \frac{z_0}{\sqrt{2\beta I}} - \frac{1}{2}, \quad (7)$$

where z_0 is the value of z which makes the integrand in Eq. (6) a maximum.

Noticing that $z_0/\sqrt{2\beta I}$ is just the average number of fermions per energy level, we can write the equation of state of the system,

$$m = \frac{1}{2\pi} \int_0^{2\pi} \frac{dk}{1 + e^{-\tilde{\varepsilon}_k(m)}} - \frac{1}{2}, \quad (8)$$

where $\tilde{\varepsilon}_k(m) = \beta J \cos k + \beta(h + 2Im)$. In the limit $T \rightarrow 0$ ($\beta \rightarrow \infty$), for $(h + 2Im)/\leq J$, Eq. (8) takes the form

$$m + \frac{1}{2} = \frac{1}{\pi} \arccos \left(-\frac{h + 2Im}{J} \right), \quad (9)$$

which, for $I = 0$, readily reduces to the well-known expression for the XX chain [3]. To analyze the behavior of the model near the quantum critical point, assuming $h \geq 0$, we define the order parameter [4] $\sigma \equiv \frac{1}{2} - m$ and expand Eq. (9) to second order in $\sigma \rightarrow 0^+$, obtaining

$$\frac{\pi^2}{2} \sigma^2 - \frac{2I}{J} \sigma = \frac{h_c - h}{J} \geq 0, \quad (10)$$

where $h_c = J - I$. For $I = 0$ we regain the usual XX chain result

$$\sigma \sim (h_c - h)^{1/2}, \quad (11)$$

while for $I < 0$ we get the expected mean-field scaling form

$$\sigma \sim (h_c - h)^1. \quad (12)$$

Note that Eq. (10) cannot be satisfied for $I > 0$, an indication that in this case the model undergoes a first-order transition at $h = h_c$ to a state where the transverse magnetization is saturated ($m = 1/2$). In this case, there is a hysteresis cycle associated to the transition, which is due to the presence of the metastable states. These states can be identified by looking at the free energy which, for $(h + 2Im)/J \leq 1$, and in the limit $T \rightarrow 0$, is explicitly given by

$$f = \frac{h}{2} - \frac{J}{\pi} \left[\sin \varphi + \left(\frac{h + 2Im}{J} \right) \varphi \right] + I(m^2 + m), \quad (13)$$

where φ is defined as

$$\varphi = \arccos \left(-\frac{h + 2Im}{J} \right). \quad (14)$$

Taking the limit $h \rightarrow 0$ in Eq. (13), and by imposing that $f(0) = f(1/2)$, which are minima of the free energy, we can show that the system presents spontaneous magnetization for $I/J \geq 4/\pi$.

The previous analysis allows us to determine the phase diagram of the model at zero temperature, shown in Figure 1. Notice that there must be a finite temperature critical line ending at the point $(h/J, I/J) = (1, 0)$, which is thus analogous to a bicritical point. The finite temperature behavior of the model will be considered in future work.

Acknowledgments. This work was partially financed by the Brazilian agencies CNPq, FINEP and Fapesp. A. P. Vieira thanks T. A. S. Haddad and S. R. Salinas for useful discussions.

References

- [1] P. Jordan, E. Wigner, Z. Physik 47 (1928) 631.
- [2] E. Lieb, T. Schultz, D. Mattis, Ann. Phys. 16 (1961) 407.
- [3] Th. Niemeijer, Physica 36 (1967) 377.
- [4] J. P. de Lima, L. L. Gonçalves, Mod. Phys. Lett. B 8 (1994) 871.

Figure caption.

Figure 1. Phase diagram of the model at $T = 0$. The solid and dashed lines indicate second- and first-order phase transitions, respectively. The diagram has of course mirror symmetry with respect to the I/J axis.

Figure 1. Goncalves et al. Abstract designation: 1S-21

